

***Testing Three Methods of Introducing Russian
Queens into Italian Honeybee Packages
SARE project FNE04-506***

Final Report submitted by:
Daniel Conlon – Project Coordinator
William Coli PhD. – Extension Specialist

Warm Colors Apiary
2 South Mill River Rd.
South Deerfield, MA. 01373

warmcolors@verizon.net 413-665-4513 www.warmcolorsapiary.com

Goals of the Project:

The goal of SARE project FNE04-506 is two-fold:

One goal is to support plans to expand Warm Colors Apiary. To do this we are increasing the number of honeybee colonies being managed for honey production, as we improve our ability to over winter colonies, and reduce the cost of maintaining our apiaries. By increasing honey production, while reducing the costs of treatments and replacement of lost colonies, we increase income and make Warm Colors Apiary a more profitable farming business.

Our second goal is to establish a sustainable population of Russian bees that are acclimated to our region. This is a key to our plans for future growth. Russian honeybees are known for their disease and mite resistance, they have a higher rate of survival in New England winters, and they produce honey crops comparable to other races of honeybee. Establishing new colonies of Russian bees can be difficult for several reasons. Russian Queens are not always accepted by other races of honeybee (Italians), comb building can be slow, and swarming is common during the first season of development.

The objectives of this project support our goals by providing practical information that can be used to improve our understanding of Russian honeybee colonies. This information will help all Northeast beekeepers improve their procedures for managing Russian bees.

Objectives:

- 1) To introduce 45 Russian Queens into Italian honeybee populations using different criteria for introduction.
 - Identify successful methods for introducing Russian Queens into Italian packages and Nucleus colonies.
- 2) Observe and record colony development.
 - Establish a baseline for package and Nucleus colony development.

- From April 2004 through April 2005.
- 3) Evaluate pre-swarm conditions and the Russian honeybees' tendency toward swarming, or early Queen Supersedure.
 - Note appearance of Queen cups and cells. Swarm cells or supercedure cells.
 - Force swarm conditions in Nucleus colonies by creating congestion in brood nest.
 - 4) Evaluate the Russian honeybee's ability to over winter by measuring strength and health of spring colonies.
 - To be completed late March or early April once weather has warmed and conditions allow, frame by frame, inspection of surviving colonies.

Participants:

Beekeepers; Jon Parrott, Kate Patterson and Susan Goddard assisted with the project by managing the test colonies, making the scheduled inspections and completing the data sheets. The forty-five colonies used in the project were located in five different apiaries. Four in Franklin County and one in Hampshire County, Massachusetts.

The collaborating beekeepers setup the hives, introduced the package bees and helped the project coordinator introduce the Russian queens. Once the queens were introduced the beekeepers fed the colonies (sugar syrup and pollen substitute), observed the scheduled events, and recorded their observations. We were able to complete the observations and record the data as planned. Our final inspections took place in late April, once the weather had warmed sufficiently to allow the hives to be opened. All surviving colonies were checked, frame by frame, to assess the colony's strength, health and condition of its Queen.

We have continued to monitor the surviving colonies through the 2005 season and collect additional information on second season development, honey production, and swarming. This additional information will appear as an update on the www.warmcolorsapiary.com website.

Project Activities:

Our desired package installation date, originally scheduled for April, was delayed due to poor weather conditions in Georgia. Dan Conlon loaded and drove the packages from Georgia to minimize the length of time they were caged. This reduced the stress to the bees and very few bees died during the trip. The packages arrived May 16th and were installed on May 16th, 17th and 18th in equipment setup in the five apiary locations. We began our test by establishing thirty new colonies using three pound packages of Italian bees purchased from Hardeman Apiaries in Mount Vernon, GA.

In addition to the package colonies we setup fifteen Nucleus ("nuc") colonies using brood and bees from over wintered Italian colonies provided by Warm Colors Apiary. Italian

queens were kept caged in each colony and Nuc to maintain morale and continuity until the Russian queens were introduced.

The package bees were placed in a single hive body using ten frames with wax foundation (duragilt). The nucs contained two frames with capped brood, two frames with honey, pollen and the adhering bees. A fifth frame with wax foundation completed each nuc. All forty five test colonies were fed syrup (1 part granulated sugar to 2 part water) continuously and given patties of pollen substitute (Bee Pro) prior to and throughout the Russian queen introduction period.

Objective #1–Introduction of 45 Russian Queens.

The Russian queens were obtained from two different breeders (suggested by Tom Rinderer of the USDA ARS), marked by Dan Conlon, the project leader, (white, green or red) and then introduced to the test colonies. The age of the Queens was three to four weeks old when introduced.

Note: Research completed by John Rhodes and Doug Somersville (May 2003) showed that successful introduction increased as Queens aged. Italian Queens that were 21-28 days of age had a 90% acceptance rate. Queen age and higher pheromone levels are thought to contribute to this improved acceptance rate.

Our objective during the introduction phase of the project was to identify conditions that would improve acceptance of Russian queens being introduced into Italian packages. The problem has been that a high percentage of Russian queens have been rejected, or superceded, (replaced by workers) during the introduction period and during the first season. Although a common complaint by beekeepers, it is not a problem for all beekeepers. This led us to consider the method of introduction plays a large part in the overall success of re-queening or starting new colonies.

Note; Queens used in this project were obtained from Hardeman Apiaries GA, Brachman Apiaries NY, and Warm Colors Apiaries MA. All were mated using Russian stock but purity of mating is not known. They are considered hybrids as they had opportunity to mate with drones of unknown origin.

We were interested in identifying procedures that helped Queens to be accepted and not necessarily the reasons they were rejected by an established colony. Common practices, known to improve a Queen's acceptance during introduction were used for all test colonies.

These included;

- Feeding colonies sugar syrup to simulate a nectar flow.
- Providing a constant supply of pollen substitute to encourage egg laying.
- Removing attendant workers from introduction cages.
- Using older mated Queens.
- Delayed release of Queen to allow workers to distribute queen substance.

We considered the introduction successful once capped brood appeared in a normal pattern.

Our test was to vary the length of time the queens were kept in cages before being released into an Italian worker population. This would allow varying intervals of time for workers to obtain and distribute queen substance to the hive population. It would also allow memory of the previous Italian queen's presence to fade. Three groups of 15 colonies were used:

Group #1 - Queens released after two days in cage (white).

Group #2 - Queens released after five days in cage (green).

Group #3 - Nucleus colonies from over wintered Italian colonies (Red). Queens were released direct into Nuc's after 24 hours.

First we removed the caged Italian Queens from the colonies. The Italian Queens kept colony morale and organization intact while the Russian queens were being shipped. This also simulated typical conditions for re-queening of a colony. Re-queening involves removing an established Queen, and then introducing her replacement.

All our Russian Queens were introduced in cages with workers (attendants) removed prior to introduction. It is well known that attendant workers may become defensive and release alarm pheromone. This creates fighting and Queens can be killed, or injured, during this behavior. By removing the attendants we eliminated one additional condition that may have caused colony workers to reject a new Queen.

Note: Russian Queens were held in a "Queen bank" before being introduced to colonies. This was necessary in order to have all queens available for the test on the same dates. We were careful to limit the number of days the queens were confined in queen banks (five days) as it may contribute to delays or problems with egg laying. A Queen bank is a Queen-less colony containing young emerging worker bees. Six to ten caged Queens (without attendant workers in cages) can be kept for short periods using this method. The young workers will feed and care for all the Queens as they have not developed an attraction to any specific Queen substance.

Colonies were not smoked, but sprayed with sugar syrup before the Queen cages were placed over the clusters. All colonies were being fed syrup & pollen substitute before the Russian Queens were introduced. We had decided not to use smoke as it can be disruptive, and may lead to workers becoming agitated and defensive. Spraying syrup has a calming effect on the bees and distracts the older foraging bees that tend to be most defensive with new Queens.

Project Timeline:

- 1) May 16th, 17th & 18th - 30 Packages (15 group #1 and 15 group #2) installed in Five Apiaries. Each location had an equal number of Group #1 and Group #2. All

15 Nucs were setup at Warm Colors Apiary. Total of 45 test colonies were started for this project. Italian Queens were kept caged to maintain colony moral.

Apiary Locations:

- Red Hen Farm – Florence, Massachusetts - four colonies.
- University of Massachusetts Agronomy Farm – six colonies.
- Bar Farm – Deerfield, Massachusetts - six colonies.
- Urkiels Farm – Whately, Massachusetts - six colonies.
- Warm Colors Apiary – South Deerfield, MA – eight colonies & 15 Nucs.


- 2) May 16th through May 20th - Russian Queens were introduced to test colonies.
- 3) May 19th through May 22nd – Group #1 Queens released from cages.
- 4) May 23rd through May 25th – Group #2 Queens released from cages.
- 5) May 16th through May 18th – Group #3 Queens released direct (after two hours in cage) into Nucs.
- 6) May – eggs, larva and capped brood observed (see attached chart for specific dates). Varied from 2 to 15 days after Queens’s release.
- 7) June & July – second hive bodies added. Varied from 30 to 98 days after Queens’s release.
- 8) July 4th – lost two test colonies to black bear at Warm Colors Apiary.
- 9) August 18th – lost six test colonies to black bear at Warm Colors Apiary.
- 10) October – Test colonies entering winter season: Group #1 – 14, Group #2 – 11, Group #3 – 12. A total of 34 test colonies remaining to begin winter.
- 11) December & January – visual examination on warm days indicted activity in all test colonies. All were active in mid-January.
- 12) Spring inspections were completed in late April of 2005. All 27 surviving colonies and nucs were active in May. Chalk brood was found in six colonies (May was wet and cool).
- 13) June all 27 surviving colonies successfully increased populations and all produced some surplus honey.
- 14) July testing found Varroa mite populations low. No treatments were used on test colonies. Screened bottom boards were in use May through August.
- 15) Final Report completed in August 2005.
- 16) Management Guide for raising Russian Honeybees completed in September 2005.

Results: Refer to the charts and tables attached to this report. They show dates and days mentioned in the following discussion of results. Complete tables including data collected are attached to this report.

Objective #1 - *The introduction of Russian Queens to Italian colonies showed no significant difference in acceptance as a result of varying release times.*

- Of the three test groups one Queen from each group was not accepted. Two after release into the colony and one was found dead in her cage. This is a 6 1/2% rejection rate for each of the three test groups. We reintroduced a second Queen to each of these colonies and were successful on the second introduction.

(See Events Table A for specific dates and days)



- All three Queens were from the same breeder.
- Specific reasons that the Queens were rejected were not determined.

- *There was no significant difference, before egg laying began, among test groups.*
 - Group #1 (Queens released from cages after 2 days) averaged 4 days.
 - Group #2 (released after 5 days) averaged 3 days.
 - Group #3 Nucs (released after 24 hours) 4 days.
- *Age of Queens may have improved acceptance.* All introduced queens were 21 to 28 days old. This is considered to be the optimal age for Queen Pheromone (Queen substance) production and improves their attractiveness to workers.
 - Queen substance is a pheromone (9ODA plus seventeen other compounds) secreted by the “mandibular” gland in the Queen, collected and distributed throughout the colony by workers.
 - High levels of Queen substance stimulates foraging, brood rearing, inhibits replacement of queens, stabilizes swarms, and acts as a sexual attractant during mating.
 - Queens produce QS at six days of age (enough to attract drones during mating flights). After three weeks mature Queens (mated) produce double the amount and do so daily.

The overall time, before eggs were found, for all groups averaged 3 to 4 days after the queen was released from her cage. This was surprising as beekeepers have complained that Russian Queens will take up to two weeks before they lay eggs. We found no difference between Russian and Italian Queens.


- *Check back time* to confirm a Queen is present and laying eggs could be set at 5 days. It is generally recommended that Italian Queens be checked one week after introduction for eggs. Based on our results we recommend Russian Queens be checked after one week. Queens taking longer to begin laying eggs should be monitored and replaced if they take longer than ten days.
 - This may be the result of mated Queens being confined in queen banks for long periods of time. It is common for breeders to hold Queens in this manner before shipping. The length of confinement can influence a queen’s egg laying ability.

Objective #2 – The rate of colony buildup varied among the individual colonies and test groups. We measured the development by counting the number of frames with drawn comb. Honeybees are limited by a lack of comb to store food and raise brood. We were also measuring the comb production by Italian workers during the initial five to six weeks, or until new Russian workers had emerged and the Italian workers had died off. This did not allow us to compare actual rate of comb production between Russian and Italian honeybees. It did provide a baseline for packages started on foundation (using 3 pound Italian packages).

The conversion from Italian workers to Russian workers required four to seven weeks. It was possible to determine the proportion of Russian to Italian workers by the color shift on the combs (golden Italians to darker Russians). All colonies were fully Russian after the eighth week. We added a 2nd hive body, after four to eight weeks, when eight of ten frames were drawn in the 1st hive body. These are typically the visual points for beekeepers to evaluate colony strength and progress.

Days to adding the Second hive body after the queen was released showed the widest difference in our test. (Refer to Events Table A & Winter Comparison Table B).

- Average number days for all test groups = 45.
- Group #1 = 59 days.
- Group #2 = 43 days.
- Group #3 = 34 days.



We had predicted that Group #1 would develop faster than #2 as the Queens were released three days earlier. It was surprising to see #1 having a higher average number of days. The release time had little to do with development. More significant was the amount of drawn comb workers had available for brood rearing.

- *Nucs had the advantage of starting with capped brood and four frames of drawn comb requiring 9 to 25 fewer days to move into a 2nd hive body.*
 - Russian colony development can be accelerated by starting packages with four or more frames of drawn comb, and at least one frame containing capped brood.

- Nucs were also started with honey & pollen. This is also a factor in rate of development.
- *Population growth, race of bees, and comb production.* Groups #1 & #2 were started with 3 pounds of package of Italian worker bees (approximately 10,500). The age of the workers, and delay in raising new workers has some impact on a colony's ability to draw new comb and maintain a larger brood nest. Our observations did not give us specific information to measure this but it was a factor when combined with below average temperatures and wet weather during the early buildup period.
 - During the Race conversion (Italian to Russian) the rate of new comb production did not appear to increase until weeks eight and nine. This was also the point when the population was mostly newly emerged Russian workers at the peak of their wax producing abilities. During the August nectar flow most of the Russian colonies finished drawing comb in both hive bodies, and one medium honey super. Thirty colonies produced surplus honey before winter.
- *The average age of worker bees is 3 to 6 weeks during the summer.* If all workers were less than ten days of age when installed in the hive, that would leave a life span of five more weeks for the Italian workers.
 - It requires 21 days for a worker to develop from an egg and emerge as an adult.
 - Three weeks after the Queen begins to lay eggs the first new replacement workers begin to emerge. This is also the earliest appearance of Russian workers from the introduced Queen.

Note: Natural swarms created to propagate new colonies are made up of 70% workers less than 10 days of age. This gives the new colony the maximum life span of its work force to draw new comb and raise replacement workers before the population dwindles below a sustainable level. Package bees are artificial swarms shaken from large colonies into cages. The age of package bees is likely to have a greater age variation and if older a tendency to dwindle in a shortened period of time.

- *Dwindling (loss of bees) of package bees was rapid.* Many of the colonies lost an estimated half of their population before the first new brood emerged.
 - Loss of workers and cool weather kept brood the nest size small limiting the rate of population growth after three weeks.
 - Samples were tested for tracheal mites but levels were very low.
 - At the UMass Agronomy Farm end hives increased due to drifting of foraging bees leaving middle colonies with fewer bees. This was the result of windy conditions during the first week, and colonies being too close during package installation.
- *Russian populations increased well during August into September.* This coincided with a strong fall nectar and pollen supply.

Objective #3 – Tendency of Russians to swarm or supercede. During the Queen introduction and race conversion phases of the project, we lost no swarms but did observe swarm cups and swarm cells. Swarming began in weeks seven and eight, after Russian workers became dominant in the colony, with many Queen cups being found in our test colonies. All Queens were marked and we could easily verify any lost Queen due to swarming or supercedure.

Note: We did not prevent the test colonies from swarming. Interference would have changed the findings that we hoped would provide a baseline for future management. We intended to allow them to develop without swarm prevention or intervention. Our only prevention was the additional space created as we added a 2nd hive body or honey super.


A total of twelve colonies swarmed during our project. Six colonies swarmed in their seven (2) and eighth (4) weeks. Six swarmed after twelve weeks.

- Queen cups appeared in a few colonies after week three and were commonly found in colonies after week nine.
- The incidence of Queen cups and Swarm cells increased as the population converted from Italian to Russian workers.
- Russian bees do produce Queen cups and remove them without actually swarming. This can be confusing to beekeepers as it does not necessarily indicate a colony intends to swarm.
- Swarming was most common with colonies and nucs that had drawn most of the comb in a given space. Early swarms appeared before, or immediately after adding the 2nd super with eight or more frames drawn. This also occurred in weeks seven and eight when Russian bees had become the dominant race.
 - In all swarming colonies the hive bodies had several frames of foundation.
 - Comb space not foundation appears to be a factor in swarming.
- Colonies that were given a 2nd hive body containing seven frames of drawn comb, and nucs that were transferred to ten frame bodies with five frames of foundation before swarm cups appeared did not swarm or swarmed late in August.
- We did not have any actual supercedure of Queens. We did see numerous Queen cups on the upper sides of frames indicating supercedure, but all marked Queens were accounted for and were not replaced.

Supercedure did not occur in our test colonies. This was unexpected as beekeepers had stated supercedure was common with Russian Queens. Supercedure happens when workers sense a Queen is failing, is injured or is otherwise unable to continue egg laying. This may indicate a problem with specific breeding stock or methods of queen rearing by some commercial breeders.

- *If their queens are not mature, healthy, and properly mated this would result in early supercedure.*

- The pressure of providing large numbers of queens early in the season pushes commercial breeders to select queens when eggs appear in the mating nucs. Queens are often two weeks of age when shipped with package bees.
- The visual confirmation of eggs does not provide sufficient information to ascertain the quality of the queen or how successfully she was inseminated during mating flights.
- Lack of sufficient drones in large breeding operations combined with poor weather conditions can reduce the number of successfully mated queens.
- *Our Queens were acquired from breeders known for producing high quality Queens and having access to the best Russian stock.*
- *The other consideration is that new lines of selected Russian stock are being released each year to commercial Queen Breeders by the USDA ARS research labs.*
 - Although the original breeding stock was found in the Primorski Peninsula of Russia, USDA researchers have been intensely selecting for desirable behaviors at the Baton Rouge USDA ARS Laboratory.
 - This ongoing selection is a breeding program to improve the strong cultural behaviors found in the Russian honeybee.
 - The Russian bee should *not* be considered a new species or new race of honeybee, but the result of a program to identify and selectively breed for mite tolerance, disease resistance, and other behaviors sought after by researchers and beekeepers.
- *The USDA ARS test laboratory, in Baton Rouge, selects to improve breeding stock and reduce undesirable behaviors in each new release. There are now at least a dozen lines of Russian stock being used in commercial breeding programs. Queens used in our tests may reflect some of these improvements.*
- *Our Russian Queens are considered to be “hybrids”. Queens are considered hybrids when mated with drones from other honeybee races; for example Italians. The purity (fully Russian) of these Queens can vary and the resultant behavior may be influenced, in part, by the drone’s genetic contribution.*



Objective #4 - Over wintering survival was about 79%. We entered winter with 34 (of the original 45) colonies in September 2004, with 27 surviving into the summer of 2005.

- 23 Colonies (Groups #1 & #2) were over wintered in two deep hive bodies with 60 – 80 pounds of honey & pollen. Colony strength and overall

size was comparable in both groups in September. 19 of the 23 survived into summer.

- 11 Nucs were returned to five frames (four full frames with honey and a center frames with partial brood). 8 of the 11 survived into summer.
- Survival rates were similar for each group: 77 to 80%. Losses were about 20%.
- This is an above average rate of loss when compared to regional losses averaging 40 to 70%.

It is difficult to know with certainty the actual reason a colony died during winter. Generally starvation, loss of the queen, and stress resulting from disease and/or mites are the primary reasons for winter loss. We knew the colonies were in good health, knew the quantities of available honey & pollen, and knew the strength of the worker population. All colonies and nucs began winter in a queen-right condition. We were able to arrive at several useful conclusions regarding Russians and their ability to winter.

- Russians use far less honey & pollen during the winter months.
- It is less likely that a Russian colony will die of starvation providing honey is available in the hive. They stay in contact with their honey reserves during long periods of cold weather.
- Russian colonies with 8 frames of honey & pollen and at least 6 frames of bees in March were strong enough to be divided in April or May.
- Spring buildup is slow until natural pollen and nectar become available.

Conditions Affecting Our Project Results:

Weather during the early test period was mild and dry. It did not have an effect on our introductions. Spring and summer were below average temperature with higher than normal rainfall. Cool wet weather may have slowed comb building and resulted in delaying colony growth. The cool temperatures, particularly at night, require bees to generate more heat. This limits the comb area that can be maintained at the proper brood incubation temperatures (92 to 96F). The result is fewer bees being raised until a sufficient population of new bees has been raised. When we compared the test colonies to other package colonies, containing Italian Queens, most were lagging behind the Italians in comb building by 6-8 frames. Cooler wet conditions also reduced the bee's ability to forage and find adequate nectar and pollen. This may have also affected their development. We have concluded that Russians tend to raise large populations only when sufficient pollen and nectar are available, and they will self-limit brood rearing in times of low natural food supplies. Beekeepers may help this situation by providing supplemental feeding between nectar flows.

Note: During the summer of 2005 we had several extended periods of heat and high humidity. June had 18 days that reached 90F with high humidity. Russian Queens were observed to have stopped laying during these extended periods, and in some cases brood was being removed from the hive without any disease being detected. This was also observed in July when we experienced a second period of high heat and humidity.

Varroa mites were more prevalent this year (2004) than we have observed in the past few seasons. They did not appear to be a factor in our test colonies, but may have sped up their infestation in apiaries shared by other mite infested hives. We will compare overwintered test colonies, isolated from sources of mite infestation, with those sharing apiaries to determine if there is any significant difference in mite loads.

Note: Varroa mites were less a problem during the 2005 season. We conduct regular field tests (sugar rolls) to identify colonies needing mite control. In 2004 we found infestations above economic thresholds to be common in all our bee yards. In 2005 we found high infestations in only three of our fourteen yards.

Black bears are now a common problem for crop farmers and beekeepers in western Massachusetts. Black bear populations have grown each year in spite of record numbers being killed by hunters. We lost eight of our test colonies to bears. All colonies lost did not have the protection of an electric fence. We lost no colonies kept within electric fences. Colonies reported by members of county associations in 2004 were estimated to be around 150 lost or severely damaged.

Note: The Black bear population in western Massachusetts is now estimated to be as high as 3800 (Mass Fish & Wildlife). We lost a total of six colonies to bears in 2005 (outside of fencing) during May pollination of fruit orchards. We have enclosed all bee yards with electric fencing and this seems to be the best method to discourage bears from feeding on bees.

Collaborating beekeepers were not always able to stay on schedule with inspections and feeding. This should not be considered a criticism of any participant's efforts. The number and frequency of observations required during this project was demanding and it was not always possible for each participant to meet every scheduled inspection. This may have had some influence on our project results. Overall we collected the information as planned and the findings did not indicate any extreme variation that may be caused by occasional delays in inspections, feeding or incomplete reporting. The purpose of this project has been to find useful information that helps us understand and better manage Russian honeybees. Our results have provided practical ideas for improving scheduled management tasks.

Assessment:

Based on our testing and observations it appears that Russians can be introduced successfully using procedures described in this report. The next step will be to establish a program that will identify Queens to use as breeding stock. A breeding program that selects for winter survival, gentle behavior, less swarming and increased honey production will, over time, improve our stock. This will lead to a sustainable population of Russian honeybees.

Note: Africanized honeybees are now established in the Southwest (Texas to California) and have recently been found in Alabama and Florida. This is a serious threat to commercial package and Queen Producers, and their ability to maintain breeding stock without the genetic influence of Africanized bees. This further underlines the importance of establishing a permanent population of honeybees in the Northeast.

Our observations also show that Russians may take a longer period to draw comb resulting in a slower population buildup than Italian honeybees. We observed that Russian brood nests would increase when natural pollen and nectar were available, and slow when nectar flows stopped. Further testing using food to stimulate faster increase in brood rearing is needed. A study comparing food would be useful in improving the rate of colony development. Although we did not specifically compare syrup to honey, or pollen substitute to natural pollen, it was observed that the Russians increased brood nest area rapidly when natural pollen was plentiful. This may indicate that feeding honey and pollen would stimulate brood rearing. Also Russians may benefit by stimulative feeding during times when natural food sources are scarce.

Swarming is a considerable problem for Russians. Losing swarms decreases colony populations and reduces honey production. Late season swarms were a particular problem. There are many manipulations that can be used to prevent or stop swarming. A swarm prevention program that works would be a valuable tool for managing Russians.

Report respectfully submitted by,

Daniel Conlon
Warm Colors Apiary
South Deerfield, Massachusetts
November 10, 2005

warmcolors@verizon.net
www.warmcolorsapiary.com
413-665-4513